



**Fermi National Accelerator Laboratory**

**FERMILAB-Pub-00/037-T**

**Explicit  $SO(10)$  Supersymmetric Grand Unified Model for the Higgs  
and Yukawa Sectors**

Carl H. Albright

*Northern Illinois University  
DeKalb, Illinois 60115*

S.M. Barr

*University of Delaware  
Newark, Delaware 19716*

*Fermi National Accelerator Laboratory  
P.O. Box 500, Batavia, Illinois 60510*

February 2000

Submitted to *Physical Review Letters*

## **Disclaimer**

*This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.*

## **Distribution**

*Approved for public release; further dissemination unlimited.*

## **Copyright Notification**

*This manuscript has been authored by Universities Research Association, Inc. under contract No. DE-AC02-76CH03000 with the U.S. Department of Energy. The United States Government and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government Purposes.*

## Explicit $SO(10)$ Supersymmetric Grand Unified Model for the Higgs and Yukawa Sectors

Carl H. Albright<sup>1</sup>  
Department of Physics  
Northern Illinois University, DeKalb, IL 60115  
and Fermi National Accelerator Laboratory  
P.O. Box 500, Batavia, IL 60510

S.M. Barr<sup>2</sup>  
Bartol Research Institute  
University of Delaware  
Newark, DE 19716

### Abstract

A complete set of fermion and Higgs superfields is introduced with well-defined  $SO(10)$  properties and  $U(1) \times Z_2 \times Z_2$  family charges from which the Higgs and Yukawa superpotentials can be constructed. The Higgs fields solve the doublet-triplet splitting problem, while the structure of the four Dirac fermion and right-handed Majorana neutrino mass matrices derived coincide with those constructed previously from an effective operator approach. The model fits the quark and lepton mass and mixing data remarkably well with the bimaximal atmospheric and solar neutrino vacuum solutions in this simplest version.

PACS numbers: 12.15.Ff, 12.10.Dm, 12.60.Jv, 14.60.Pq

Key words: SUSY,  $SO(10)$ , quark and lepton mass matrices

---

<sup>1</sup>E-mail: albright@fnal.gov; <sup>2</sup>E-mail: smbarr@bartol.udel.edu

In a series of recent papers [1] - [4] the authors have shown how fermion mass matrices can be constructed in an  $SO(10)$  supersymmetric grand unified framework by use of a minimal Higgs structure which solves the doublet-triplet splitting problem [5]. The construction was carried out in an effective operator approach with phenomenological input, including the Georgi-Jarlskog relations [6]. Here we show how one can introduce a set of matter and Higgs  $SO(10)$  superfields with  $U(1) \times Z_2 \times Z_2$  family charges from which the derived Higgs and Yukawa superpotentials uniquely give the structure of the fermion mass matrices previously obtained. The quark and lepton mass and mixing data are reproduced remarkably well with the solar neutrino vacuum solution preferred, provided the up quark mass is not zero at the GUT scale – otherwise the small angle MSW solution [7] is obtained. The right-handed Majorana neutrino matrix arises from a Higgs field which couples pairs of superheavy conjugate neutrino singlets.

We begin with a listing in Table I. of the Higgs and matter superfields in the proposed model along with their family charges. As demonstrated in [5], in order to do all

#### Higgs Fields Needed to Solve the 2-3 Problem:

$$\begin{aligned}
\mathbf{45}_{B-L}: & \quad A(0)^{+-} \\
\mathbf{16}: & \quad C(\tfrac{3}{2})^{-+}, C'(\tfrac{3}{2} - p)^{-+} \\
\overline{\mathbf{16}}: & \quad \bar{C}(-\tfrac{3}{2})^{++}, \bar{C}'(-\tfrac{3}{2} - p)^{-+} \\
\mathbf{10}: & \quad T_1(1)^{++}, T_2(-1)^{+-} \\
\mathbf{1}: & \quad X(0)^{++}, P(p)^{+-}, Z_1(p)^{++}, Z_2(p)^{++}
\end{aligned}$$

#### Additional Higgs Fields for the Mass Matrices:

$$\begin{aligned}
\mathbf{10}: & \quad T_0(1 + p)^{+-}, T'_0(1 + 2p)^{+-}, \\
& \quad \bar{T}_0(-3 + p)^{-+}, \bar{T}'_0(-1 - 3p)^{-+} \\
\mathbf{1}: & \quad Y(2)^{-+}, Y'(2)^{++}, S(2 - 2p)^{--}, S'(2 - 3p)^{--}, \\
& \quad V_M(4 + 2p)^{++}
\end{aligned}$$

Table I. Higgs superfields in the proposed model.

the symmetry breaking, one  $\mathbf{45}$  adjoint Higgs with its VEV pointing in the  $B - L$  direction, a pair of  $\mathbf{16} + \overline{\mathbf{16}}$  spinor Higgs, plus a pair of  $\mathbf{10}$  vector Higgs and several Higgs singlets are required. In order to complete the construction of the Dirac mass matrices, four more vector Higgs and four additional singlets are needed. Finally, one Higgs singlet is introduced

to generate the right-handed Majorana mass matrix.

From the Higgs  $SO(10)$  and family assignments, it is then possible to write down explicitly the full Higgs superpotential, where we have written it as the sum of five terms:

$$\begin{aligned}
W_{\text{Higgs}} &= W_A + W_{CA} + W_{2/3} + W_{HD} + W_R \\
W_A &= \text{tr} A^4 / M + M_A \text{tr} A^2 \\
W_{CA} &= X(\overline{C}C)^2 / M_C^2 + F(X) \\
&\quad + \overline{C}'(PA/M_1 + Z_1)C + \overline{C}(PA/M_2 + Z_2)C' \\
W_{2/3} &= T_1 A T_2 + Y' T_2^2 \\
W_{HD} &= T_1 \overline{C} \overline{C} Y' / M + \overline{T}_0 C C' + \overline{T}_0 (T_0 S + T_0' S') \\
W_R &= \overline{T}_0 \overline{T}_0' V_M
\end{aligned} \tag{1}$$

The Higgs singlets are all assumed to develop VEV's at the GUT scale.  $W_A$  fixes  $\langle A \rangle$  through the  $F_A = 0$  condition where one solution is  $\langle A \rangle \propto B - L$ , the Dimopoulos-Wilczek solution [8].  $W_{CA}$  gives a GUT-scale VEV to  $\overline{C}$  and  $C$  by the  $F_X = 0$  condition and also couples the adjoint  $A$  to the spinors  $C$ ,  $\overline{C}$ ,  $C'$  and  $\overline{C}'$  without destabilizing the Dimopoulos-Wilczek solution or giving Goldstone modes.  $W_{2/3}$  gives the doublet-triplet splitting by the Dimopoulos-Wilczek mechanism.  $W_{HD}$  mixes the  $(1, 2, -1/2)$  doublet in  $T_1$  with those in  $C'$  (by  $F_{\overline{C}} = 0$ ), and in  $T_0$  and  $T_0'$  (by  $F_{\overline{T}_0} = 0$ ).

To fill out the model, we specify the  $SO(10) \times U(1) \times Z_2 \times Z_2$  quantum numbers of the various matter fields in Table II. We require three chiral spinor fields  $\mathbf{16}_i$ , one for each light family, two vector-like pairs of  $\mathbf{16} + \overline{\mathbf{16}}$  spinors which can get superheavy, a pair of superheavy  $\mathbf{10}$  fields in the vector representation, and three pairs of superheavy  $\mathbf{1} - \mathbf{1}^c$  fermion singlets.

$\mathbf{16}_1(-\frac{1}{2} - 2p)^{+-}$	$\mathbf{16}_2(-\frac{1}{2} + p)^{++}$	$\mathbf{16}_3(-\frac{1}{2})^{++}$
$\mathbf{16}(-\frac{1}{2} - p)^{-+}$	$\mathbf{16}'(-\frac{1}{2})^{-+}$	
$\overline{\mathbf{16}}(\frac{1}{2})^{+-}$	$\overline{\mathbf{16}}'(-\frac{3}{2} + 2p)^{+-}$	
$\mathbf{10}_1(-1 - p)^{-+}$	$\mathbf{10}_2(-1 + p)^{++}$	
$\mathbf{1}_1(2 + 2p)^{+-}$	$\mathbf{1}_2(2 - p)^{++}$	$\mathbf{1}_3(2)^{++}$
$\mathbf{1}_1^c(-2 - 2p)^{+-}$	$\mathbf{1}_2^c(-2)^{+-}$	$\mathbf{1}_3^c(-2 - p)^{++}$

Table II. Matter superfields in the proposed model.

In terms of these fermion fields and the Higgs fields previously introduced, one can then spell out all the terms in the Yukawa superpotential which follow from their  $SO(10)$  and  $U(1) \times Z_2 \times Z_2$  assignments:

$$\begin{aligned}
W_{Yukawa} = & \mathbf{16}_3 \cdot \mathbf{16}_3 \cdot T_1 + \mathbf{16}_2 \cdot \mathbf{16} \cdot T_1 + \mathbf{16}' \cdot \mathbf{16}' \cdot T_1 \\
& + \mathbf{16}_3 \cdot \mathbf{16}_1 \cdot T'_0 + \mathbf{16}_2 \cdot \mathbf{16}_1 \cdot T_0 + \mathbf{16}_3 \cdot \overline{\mathbf{16}} \cdot A \\
& + \mathbf{16}_1 \cdot \overline{\mathbf{16}}' \cdot Y' + \mathbf{16} \cdot \overline{\mathbf{16}} \cdot P + \mathbf{16}' \cdot \overline{\mathbf{16}}' \cdot S \\
& + \mathbf{16}_3 \cdot \mathbf{10}_2 \cdot C' + \mathbf{16}_2 \cdot \mathbf{10}_1 \cdot C + \mathbf{10}_1 \cdot \mathbf{10}_2 \cdot Y \\
& + \mathbf{16}_3 \cdot \mathbf{1}_3 \cdot \overline{C} + \mathbf{16}_2 \cdot \mathbf{1}_2 \cdot \overline{C} + \mathbf{16}_1 \cdot \mathbf{1}_1 \cdot \overline{C} \\
& + \mathbf{1}_3 \cdot \mathbf{1}_3^c \cdot Z + \mathbf{1}_2 \cdot \mathbf{1}_2^c \cdot P + \mathbf{1}_1 \cdot \mathbf{1}_1^c \cdot X \\
& + \mathbf{1}_3^c \cdot \mathbf{1}_3^c \cdot V_M + \mathbf{1}_1^c \cdot \mathbf{1}_2^c \cdot V_M
\end{aligned} \tag{2}$$

where the coupling parameters have been suppressed. To obtain the GUT scale structure for the fermion mass matrix elements, all but the three chiral spinor fields in the first line of Table II. will be integrated out. The right-handed Majorana matrix elements will all be generated through the Majorana couplings of the  $V_M$  Higgs field with conjugate singlet fermions as given above.

The procedure for deriving the Dirac mass matrices  $U$ ,  $D$ ,  $L$ , and  $N$  is the following. For each type of fermion  $f$ , where  $f = u_L, u_L^c, d_L, d_L^c, \ell_L^-, \ell_L^+, \nu_L$  and  $\nu_L^c$ , the superheavy mass matrix connecting the  $f$  to the  $SU(3) \times SU(2) \times U(1)$ -conjugate representation  $\overline{f}$  is first found from Eq. (2) by setting the weak-scale VEV's and the intermediate-scale VEV,  $V_M$ , to zero. This will give three zero mass eigenstates for each type of  $f$ , corresponding to the three light families. Then the terms in Eq. (2) involving  $\langle T_1 \rangle$ ,  $\langle C' \rangle$ ,  $\langle T_0 \rangle$ , and  $\langle T'_0 \rangle$  give rise to the  $3 \times 3$  Dirac mass matrices coupling  $u_L$  to  $u_L^c$ , etc. This procedure is spelled out explicitly in [9].

Under the assumption that the zero-mass states have their large components in the chiral representations  $\mathbf{16}_1$ ,  $\mathbf{16}_2$  and  $\mathbf{16}_3$ , and all the other components are small, the Dirac mass matrices obtained have precisely the structure previously found in our studies by means of an effective operator approach:

$$\begin{aligned}
U &= \begin{pmatrix} \eta & 0 & 0 \\ 0 & 0 & \epsilon/3 \\ 0 & -\epsilon/3 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} 0 & \delta & \delta' e^{i\phi} \\ \delta & 0 & \sigma + \epsilon/3 \\ \delta' e^{i\phi} & -\epsilon/3 & 1 \end{pmatrix}, \\
N &= \begin{pmatrix} \eta & 0 & 0 \\ 0 & 0 & -\epsilon \\ 0 & \epsilon & 1 \end{pmatrix}, \quad L = \begin{pmatrix} 0 & \delta & \delta' e^{i\phi} \\ \delta & 0 & -\epsilon \\ \delta' e^{i\phi} & \sigma + \epsilon & 1 \end{pmatrix},
\end{aligned} \tag{3}$$

with  $U$  and  $N$  in units of  $M_U$  and  $D$  and  $L$  in units of  $M_D$ . The matrix parameters are identified with the Yukawa couplings and Higgs couplings and VEV's as follows:

$$\begin{aligned}
M_U &= (t_3)_{5(10)}, & M_D &= (t_3)_{\bar{5}(10)}, \\
\epsilon M_U &= |3(a_q/p)(t_2)_{5(10)}|, & \epsilon M_D &= |3(a_q/p)(t_2)_{\bar{5}(10)}|, \\
\eta M_U &= (y'/s'')^2 (t')_{5(10)}, & \sigma M_D &= -(c/y)(c')_{\bar{5}(16)}, \\
& & \delta M_D &= t_0 \bar{t}_0 / s, \\
& & \delta' M_D &= (t'_0 \bar{t}_0 / s') e^{-i\phi},
\end{aligned} \tag{4}$$

where the subscripts on  $t_2$ ,  $t_3$ ,  $t'$  and  $c'$  refer to the  $SU(5)[SO(10)]$  representation content of the VEV's. The following shorthand notation has been introduced

$$\begin{aligned}
t_3 &= \lambda_{16_3 16_3 T_1} \langle T_1 \rangle, & t_2 &= \lambda_{16_2 16 T_1} \langle T_1 \rangle, \\
t' &= \lambda_{16' 16' T_1} \langle T_1 \rangle, & c' &= \lambda_{16_3 10_2 C'} \langle C' \rangle, \\
c &= \lambda_{16_2 10_1 C} \langle C \rangle, & \bar{c}_i &= \lambda_{16_i 1_i \bar{C}} \langle \bar{C} \rangle, i = 1, 2, 3, \\
p &= \lambda_{16 \bar{16} P} \langle P \rangle, & p_{22} &= \lambda_{1_2 1_2^c P} \langle P \rangle, \\
a_q &= \lambda_{16_3 \bar{16} A} \langle A \rangle_{B=1/3}, & x &= \lambda_{1_1 1_1^c X} \langle X \rangle, \\
y &= \lambda_{10_1 10_2 Y} \langle Y \rangle, & y' &= \lambda_{16_1 \bar{16} Y'} \langle Y' \rangle, \\
z &= \lambda_{1_3 1_3^c Z} \langle Z \rangle, & s &= \lambda_{T_0 \bar{T}_0 S} \langle S \rangle, \\
s' &= \lambda_{T'_0 \bar{T}_0 S'} \langle S' \rangle, & s'' &= \lambda_{16' \bar{16}' S} \langle S \rangle, \\
t_0 &= \lambda_{16_1 16_2 T_0}, & t'_0 &= \lambda_{16_1 16_3 T'_0}, \\
\bar{t}_0 &= \lambda_{CC' \bar{T}_0} \langle C \rangle \langle C' \rangle.
\end{aligned} \tag{5}$$

The parameter  $\eta$  is introduced to give a tiny non-zero mass to the up quark at the  $\Lambda_G$  scale. Its appearance in  $N$  will also play an important role in the determination of the type of solar neutrino solution. It should also appear in  $D$  and  $L$  but its effect is negligibly small there and of no consequence, so it is dropped. The only phase then appearing in the matrices is  $\phi$  associated with  $\delta'$ , as other phases are unphysical and can be rotated away with the exception of that associated with  $\epsilon$ . It turns out, however, that the best fits to the data

prefer a real  $\epsilon$ . Hence  $\phi$  which can be identified with the complexity of the VEV of the  $S'$  Higgs singlet is solely responsible for CP-violation in the quark sector. The structures of the matrix elements given in Eqs. (3), (4) and (5) can be understood in terms of simple Froggatt-Nielsen diagrams [10] given in [9].

Note that the 33 elements of the Dirac mass matrices are scaled by the VEV's of the  $\mathbf{10}$ ,  $T_1$ . But the  $F = 0$  conditions for the Higgs superpotential require that the pair of Higgs doublets which remain light down to the electroweak scale arise from  $5(T_1)$ ,  $\bar{5}(T_1)$ ,  $\bar{5}(C')$  and, to a very small extent from  $T_0$  and  $T'_0$  terms, which are ignored here. In particular, we can write in terms of a mixing angle  $\gamma$

$$H_U = 5(T_1), \quad H_D = \bar{5}(T_1) \cos \gamma + \bar{5}(C') \sin \gamma \quad (6)$$

whereas the orthogonal combination has become superheavy at the GUT scale. Thus the ratio of the 33 mass matrix elements found from Eqs. (4) and (6) is given in terms of the VEV's,  $v_u$  and  $v_d$  of  $H_U$  and  $H_D$ , respectively, by

$$M_U/M_D = v_u/(v_d \cos \gamma) \equiv \tan \beta / \cos \gamma \quad (7)$$

Hence we find that the large  $M_U/M_D$  ratio required for the top to bottom quark masses can be achieved with a *moderate*  $\tan \beta$  provided  $\cos \gamma$  is small.

Turning to the right-handed Majorana mass matrix, we use the zero mass left-handed conjugate states that were found implicitly above for the Dirac matrix  $N$  to form the basis for  $M_R$ . The right-handed Majorana neutrino matrix is then obtained from the last two terms in Eq. (2), and we find

$$M_R = \begin{pmatrix} 0 & A\epsilon^3 & 0 \\ A\epsilon^3 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \Lambda_R \quad (8)$$

where

$$\begin{aligned} \Lambda_R &= \lambda_{1_3^c 1_3^c V_M} \langle V_M \rangle (\bar{c}_3/z)^2, \\ A\epsilon^3 \Lambda_R &= \lambda_{1_1^c 1_2^c V_M} \langle V_M \rangle (\bar{c}_1/x)(\bar{c}_2/p_{22}) \end{aligned} \quad (9)$$

Note that the whole right-handed Majorana mass matrix has been generated in this simple model by the one Majorana VEV coupling superheavy conjugate fermion singlets. By means of the seesaw formula [11], one can then compute the light neutrino mass matrix



$$M_\nu = N^T M_R^{-1} N = \begin{pmatrix} 0 & 0 & -\frac{\eta}{A\epsilon^2} \\ 0 & \epsilon^2 & \epsilon \\ -\frac{\eta}{A\epsilon^2} & \epsilon & 1 \end{pmatrix} M_U^2 / \Lambda_R \quad (10)$$

We now address the predictions of the mass matrices. For this purpose it is convenient to replace the parameters  $\delta$  and  $\delta'$  by

$$t_L e^{i\theta} \equiv \frac{\delta - \sigma \delta' e^{i\phi}}{\sigma \epsilon / 3}, \quad t_R \equiv \frac{\delta \sqrt{\sigma^2 + 1}}{\sigma \epsilon / 3} \quad (11)$$

which are essentially the left-handed and right-handed Cabibbo angles. In terms of the dimensionless parameters  $\epsilon$ ,  $\sigma$ ,  $t_L$ ,  $t_R$ ,  $e^{i\theta}$ ,  $\eta$ ,  $A$ , and  $M_U/M_D$ , we then find at the GUT scale

$$\begin{aligned} m_t^0/m_b^0 &\cong (\sigma^2 + 1)^{-1/2} M_U/M_D, & m_u^0/m_t^0 &\cong \eta, \\ m_c^0/m_t^0 &\cong \frac{1}{9}\epsilon^2 \cdot [1 - \frac{2}{9}\epsilon^2], & m_b^0/m_\tau^0 &\cong 1 - \frac{2}{3}\frac{\sigma}{\sigma^2+1}\epsilon, \\ m_s^0/m_b^0 &\cong \frac{1}{3}\epsilon\frac{\sigma}{\sigma^2+1} \cdot [1 + \frac{1}{3}\epsilon\frac{1-\sigma^2-\sigma\epsilon/3}{\sigma(\sigma^2+1)} + \frac{1}{2}(t_L^2 + t_R^2)], \\ m_d^0/m_s^0 &\cong t_L t_R \cdot [1 - \frac{1}{3}\epsilon\frac{\sigma^2+2}{\sigma(\sigma^2+1)} - (t_L^2 + t_R^2) \\ &\quad + (t_L^4 + t_L^2 t_R^2 + t_R^4)], \\ m_\mu^0/m_\tau^0 &\cong \epsilon\frac{\sigma}{\sigma^2+1} \cdot [1 + \epsilon\frac{1-\sigma^2-\sigma\epsilon}{\sigma(\sigma^2+1)} + \frac{1}{18}(t_L^2 + t_R^2)], \\ m_e^0/m_\mu^0 &\cong \frac{1}{9}t_L t_R \cdot [1 - \epsilon\frac{\sigma^2+2}{\sigma(\sigma^2+1)} + \epsilon^2\frac{\sigma^4+9\sigma^2/2+3}{\sigma^2(\sigma^2+1)^2} \\ &\quad - \frac{1}{9}(t_L^2 + t_R^2)], \\ V_{cb}^0 &\cong \frac{1}{3}\epsilon\frac{\sigma^2}{\sigma^2+1} \cdot [1 + \frac{2}{3}\epsilon\frac{1}{\sigma(\sigma^2+1)}], \\ V_{us}^0 &\cong t_L [1 - \frac{1}{2}t_L^2 - t_R^2 + t_R^4 + \frac{5}{2}t_L^2 t_R^2 + \frac{3}{8}t_L^4 \\ &\quad - \frac{\epsilon}{3\sigma\sqrt{\sigma^2+1}}\frac{t_R}{t_L}e^{-i\theta}], \\ V_{ub}^0 &\cong \frac{1}{3}t_L\epsilon\frac{1}{\sigma^2+1}[\sqrt{\sigma^2+1}\frac{t_R}{t_L}e^{-i\theta}(1 - \frac{1}{3}\epsilon\frac{\sigma}{\sigma^2+1}) \\ &\quad - (1 - \frac{2}{3}\epsilon\frac{\sigma}{\sigma^2+1})], \\ m_2^0/m_3^0 &\cong \left(\frac{\eta}{A\epsilon\sqrt{1+\epsilon^2}}\right) \left[1 + \frac{\eta}{A\epsilon^3\sqrt{1+\epsilon^2}}\right], \\ m_1^0/m_3^0 &\cong \left(\frac{\eta}{A\epsilon\sqrt{1+\epsilon^2}}\right) \left[1 - \frac{\eta}{2A\epsilon^3\sqrt{1+\epsilon^2}}\right], \\ U_{\mu 3}^0 &\cong -\frac{1}{\sqrt{\sigma^2+1}}(\sigma - \epsilon\frac{\sigma^2}{\sigma^2+1}), \\ U_{e 2}^0 &\cong -\frac{1}{\sqrt{2}} \left[1 - \frac{\epsilon}{3\sigma}t_L e^{i\theta} \right. \\ &\quad \left. + \frac{1}{3\sqrt{\sigma^2+1}}(1 + \epsilon\sigma)t_R\right], \\ U_{e 3}^0 &\cong \frac{1}{3\sqrt{\sigma^2+1}}(\sigma - \epsilon)t_R - \frac{\eta}{A\epsilon^2} \end{aligned} \quad (12)$$

Note that the Georgi-Jarlskog relations [6],  $m_s^0 \cong \frac{1}{3}m_\mu^0$  and  $m_d^0 \cong 3m_e^0$ , emerge as required

by design. The quark and charged lepton data are best fit at the low scale (see below) by assigning the following values to the model parameters:  $\epsilon = 0.145$ ,  $\sigma = 1.78$ ,  $t_L = 0.236$ ,  $t_R = 0.205$ ,  $\theta = 34^\circ$  (corresponding to  $\delta = 0.0086$ ,  $\delta' = 0.0079$ ,  $\phi = 54^\circ$ ),  $\eta = 8 \times 10^{-6}$ , and  $M_U/M_D \simeq 113$ .

As noted earlier, in order to obtain the simple mass matrices in Eq. (3), we had to assume that the zero-mass states have their large components in the chiral representations **16**<sub>1</sub>, **16**<sub>2</sub>, and **16**<sub>3</sub>. The conditions on the state normalization factors are all satisfied provided the following ratios are much less than unity:

$$\begin{aligned} (a/p)^2, (y'/s'')^2, (c/y)^2, \\ (\bar{c}_1/x)^2, (\bar{c}_2/p_{22})^2, (\bar{c}_3/z)^2 \ll 1 \end{aligned} \quad (13)$$

With the numerical choice of parameters given above and near equality of the various Higgs couplings, we find  $(a/p)^2 \simeq 0.02$  and  $(y'/s'')^2 \sim 6 \times 10^{-6}$ , so the first two conditions are easily satisfied. Requiring that  $(c/y)^2 \ll 1$  and with the expression for  $\sigma$  obtained from Eqs. (4), we find

$$\begin{aligned} \tan \gamma &\equiv \frac{\langle \bar{5}(C') \rangle}{\langle \bar{5}(T_1) \rangle} \gg \sigma \\ \tan \beta &\simeq \sqrt{\sigma^2 + 1} (\cos \gamma) m_t^0 / m_b^0 \ll m_t^0 / m_b^0 \end{aligned} \quad (14)$$

in terms of the  $T_1 - C'$  mixing angle,  $\gamma$ , in Eq. (6). With  $c/y \cong 0.1$ , for example,  $\tan \gamma \simeq 18$  which implies  $\tan \beta \simeq 6$ , a very reasonable mid-range value allowed by experiment. The others can also be satisfied [9].

In [9] we have evolved the results in Eqs. (12) down to the low scales with a value for  $\tan \beta = 5$ ,  $\Lambda_G = 2 \times 10^{16}$  GeV,  $\Lambda_{SUSY} = m_t(m_t)$ ,  $\alpha_s(M_Z) = 0.118$ ,  $\alpha(M_Z) = 1/127.9$ , and  $\sin^2 \theta_W = 0.2315$ . With the quantities  $m_t(m_t) = 165$  GeV,  $m_\tau = 1.777$  GeV,  $m_\mu = 105.7$  MeV,  $m_e = 0.511$  MeV,  $m_u = 4.5$  MeV,  $V_{us} = 0.220$ ,  $V_{cb} = 0.0395$ , and  $\delta_{CP} = 64^\circ$  used to determine the input parameters,  $M_U \simeq 113$  GeV,  $M_D \simeq 1$  GeV, and  $\sigma$ ,  $\epsilon$ ,  $t_L$ ,  $t_R$ ,  $\theta$  and  $\eta$  given earlier, the following values are obtained compared with experiment [12] in parentheses:

$$\begin{aligned} m_c(m_c) &= 1.23 \text{ GeV} & (1.27 \pm 0.1 \text{ GeV}) \\ m_b(m_b) &= 4.25 \text{ GeV} & (4.26 \pm 0.11 \text{ GeV}) \\ m_s(1 \text{ GeV}) &= 148 \text{ MeV} & (175 \pm 50 \text{ MeV}) \\ m_d(1 \text{ GeV}) &= 7.9 \text{ MeV} & (8.9 \pm 2.6 \text{ MeV}) \\ |V_{ub}/V_{cb}| &= 0.080 & (0.090 \pm 0.008) \end{aligned} \quad (15)$$

where the finite SUSY loop corrections for  $m_b$  and  $m_s$  have been scaled to give  $m_b(m_b) \simeq 4.25$  GeV for  $\tan \beta = 5$ .

The effective light neutrino mass matrix of Eq. (10) leads to bimaximal mixing with a large angle solution for atmospheric neutrino oscillations [13] and the “just-so” vacuum solution [14] involving two pseudo-Dirac neutrinos, if we set  $\Lambda_R = 2.4 \times 10^{14}$  GeV and  $A = 0.05$ . We then find

$$\begin{aligned}
m_3 &= 54.3 \text{ meV}, \quad m_2 = 59.6 \text{ } \mu\text{eV}, \quad m_1 = 56.5 \text{ } \mu\text{eV} \\
U_{e2} &= 0.733, \quad U_{e3} = 0.047, \quad U_{\mu 3} = -0.818, \quad \delta'_{CP} = -0.2^\circ \\
\Delta m_{23}^2 &= 3.0 \times 10^{-3} \text{ eV}^2, \quad \sin^2 2\theta_{atm} = 0.89 \\
\Delta m_{12}^2 &= 3.6 \times 10^{-10} \text{ eV}^2, \quad \sin^2 2\theta_{solar} = 0.99 \\
\Delta m_{13}^2 &= 3.0 \times 10^{-3} \text{ eV}^2, \quad \sin^2 2\theta_{reac} = 0.009
\end{aligned} \tag{16}$$

The effective scale of the right-handed Majorana mass contribution occurs two orders of magnitude lower than the SUSY GUT scale of  $\Lambda_G = 1.2 \times 10^{16}$  GeV. The effective two-component reactor mixing angle given above should be observable at a future neutrino factory, whereas the present limit from the CHOOZ experiment [15] is approximately 0.1 for the above  $\Delta m_{23}^2$ . In principle, the parameter  $A$  appearing in  $M_R$  can also be complex, but we find that in no case does the leptonic CP-violating phase,  $\delta'_{CP}$  exceed  $10^\circ$  in magnitude. Hence the model predicts leptonic CP-violation will be unobservable.

The vacuum solar solution depends critically on the appearance of the parameter  $\eta$  in the matrix  $N$ , corresponding to the non-zero  $\eta$  entry in  $U$  which gives the up quark a mass at the GUT scale. Should we set  $\eta = 0$ , only the small-angle MSW solution [7] would be obtained for the solar neutrino oscillations. The large angle MSW solution is disfavored by the larger hierarchy, i.e., very small  $A$  value, required in  $M_R$ .

In summary, we have constructed an explicit  $SO(10)$  supersymmetric grand unified model for the Higgs and Yukawa superpotentials which reproduces the fermion mass matrices previously obtained in an effective operator approach. All the quark and lepton mass and mixing data are fit remarkably well with a  $\tan \beta$  in the range of 5 - 10 with matrix parameters which are also quite reasonable.

The research of SMB was supported in part by the Department of Energy under contract No. DE-FG02-91ER-40626. Fermilab is operated by Universities Research Association Inc. under contract No. DE-AC02-76CH03000 with the Department of Energy.

## References

- [1] C.H. Albright and S.M. Barr, Phys. Rev. D **58**, 013002 (1998).
- [2] C.H. Albright, K.S. Babu, and S.M. Barr, Phys. Rev. Lett. **81**, 1167 (1998); C.H. Albright, K.S. Babu, and S.M. Barr, Nucl. Phys. B (Proc. Suppl.) **77**, 308 (1999).
- [3] C.H. Albright and S.M. Barr, Phys. Lett. B **452**, 287 (1999).
- [4] C.H. Albright and S.M. Barr, Phys. Lett. B **461**, 218 (1999).
- [5] S.M. Barr and S. Raby, *Phys. Rev. Lett.* **79**, 4748 (1997).
- [6] H. Georgi and C. Jarlskog, *Phys. Lett.* **B86**, 297 (1979); A. Kusenko and R. Shrock, *Phys. Rev.* **D49**, 4962 (1994).
- [7] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978); S.P. Mikheyev and A. Yu. Smirnov, *Yad. Fiz.* **42**, 1441 (1985), [*Sov. J. Nucl. Phys.* **42**, 913 (1985)].
- [8] S. Dimopoulos and F. Wilczek, report No. NSF-ITP-82-07 (1981), in *The unity of fundamental interactions*, Proceedings of the 19th Course of the International School of Subnuclear Physics, Erice, Italy, 1981, ed. A. Zichichi (Plenum Press, New York, 1983); K.S. Babu and S.M. Barr, Phys. Rev. D **48**, 5354 (1993).
- [9] C.H. Albright and S.M. Barr, long paper in preparation.
- [10] C.D. Froggatt and H.B. Nielsen, Nucl. Phys. B **147**, 277 (1979).
- [11] M. Gell-Mann, P. Ramond, and R. Slansky, Report No. CALT-68-709: *Supergravity* (North Holland, Amsterdam, 1979); T. Yanagida, in Proceedings of the Workshop on Unified Theory and Baryon Number of the Universe, KEK, Japan, 1979 (unpublished).
- [12] *Table of Particle Properties*, Particle Data Group, (1998); M. Bargiotti et al., hep-ph/0001293.
- [13] Y. Fukuda et al., Phys. Rev. Lett. **81**, 1562 (1998); Y. Suzuki, in Proceedings of the WIN-99 Workshop, Cape Town, 25 - 30 January 1999, to be published.

- [14] H. Fritzsch and Z.Z. Xing, Phys. Lett. B **372**, 265 (1996); E. Torrente-Lujan, Phys. Lett. B **389**, 557 (1996); V. Barger, P. Pakvasa, T.J. Weiler, and K. Whisnant, Phys. Lett. B **437**, 107 (1998); A. Baltz, A.S. Goldhaber and M. Goldhaber, Phys. Rev. Lett. **58**, 5730 (1998); H. Georgi and S.L. Glashow, hep-ph/9808293; C. Giunti, Phys. Rev. D **59**, 077301 (1999).
- [15] CHOOZ Collab. (M. Apollonio et al.), Phys. Lett. B **420**, 397 (1998).